

**EMBARGOED FOR RELEASE until April 15, 2010 – 5:00 PM (EDT)**

**ASN Contact:** Shari Leventhal • 202-416-0658 (p) • sleventhal@asn-online.org

## **WEEKEND HOSPITAL ADMISSIONS ARE HIGHER RISK FOR PATIENTS WITH ACUTE KIDNEY INJURY**

**Acute Kidney Injury Patients are Less Likely to Die when Admitted on a Weekday**

**Washington, DC (April 12, 2010)** — Patients with acute kidney injury (AKI) who are admitted to the hospital on a weekend are more likely to die than those admitted on a weekday, according to a study appearing in an upcoming issue of the *Journal of the American Society Nephrology* (JASN). This disparity was most evident in smaller hospitals. The findings indicate that researchers should further investigate the availability and timing of care to patients hospitalized with AKI.

Studies have shown that for a variety of acute illnesses, patients admitted to the hospital on a weekend are more likely to die than those admitted on a weekday. To investigate whether this is true for patients with AKI, Matthew James, MD (University of Calgary, in Alberta, Canada), Glenn Chertow, MD (Stanford University School of Medicine), and colleagues examined information from the Nationwide Inpatient Sample, a large database of admissions to acute care, nonfederal hospitals in the United States. They identified 963,730 admissions with a diagnosis of AKI between 2003 and 2006. Of these, 214,962 admissions (22%) designated AKI as the primary reason for admission (45,203 on a weekend and 169,759 on a weekday).

The investigators found that during hospitalization, 14,686 (6.8%) patients admitted with a primary diagnosis of AKI died. Patients admitted with a primary diagnosis of AKI on a weekend had a 22% increased risk of dying by day three of admission, and a 7% increase of dying during the duration of their hospital stay, compared with patients admitted on a weekday. In small hospitals, the risk was even greater. There, compared with patients admitted on a weekday, a patient's risk of dying after admission on a weekend for AKI was 34% higher after day three of admission and 17% higher over the duration of their hospital stay. The investigators also found an increased risk of death following a weekend admission among patients who were admitted for other conditions but later diagnosed with AKI during their hospital stay.

"There are many reasons why this research is important," said Chertow. "Optimizing patient safety and nephrology resource utilization are among the two most relevant," he noted.

While the underlying reasons for the observed increased deaths among AKI patients admitted on a weekend are unknown and require further investigation, other studies

have described delays in assessment, diagnosis, and management of other acute medical conditions on weekends.

Study co-authors include Ron Wald, MDCM, Chaim M. Bell, MD, PhD, FRCPC (University of Toronto, in Ontario, Canada); Marcello Tonelli, MD (University of Alberta, in Alberta, Canada), Brenda R. Hemmelgarn, MD, PhD, FRCPC (University of Calgary, in Alberta, Canada); and Sushrut S. Waikar, MD (Harvard Medical School).

In reviewing the results of Dr. James' and Dr. Chertow's study in an accompanying editorial, William McClellan, MD (Emory University) stated that the study was well-designed and that its results should encourage investigators to identify "potentially modifiable risk factors that contribute to mortality differences" so that efforts can be made to reduce AKI patients' risk of dying when admitted to the hospital on a weekend. He noted that current guidelines recommend that a patient who is admitted to the hospital for AKI should receive a timely consultation with a kidney specialist, a determination of the cause and severity of his or her condition, appropriate medications and nutritional support, and various other attributes of care. According to Dr. McClellan, it may be informative to determine how differences in these and other aspects of care might contribute to weekday-to-weekend and hospital-to-hospital variability in mortality of patients with AKI.

The article, entitled "Weekend Hospital Admission, Acute Kidney Injury, and Mortality," (doi 10.1681/ASN.2009070682) and editorial entitled "Variations in Mortality among Hospitalizations for Acute Kidney Injury," (doi 10.1681/ASN.2010030300) will appear online at <http://jasn.asnjournals.org/> on April 15, 2010.

Disclosures: The authors of the study and editorial reported no financial disclosures.

*The American Society of Nephrology (ASN) does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies. The content of this article does not reflect the views or opinions of ASN. Responsibility for the information and views expressed therein lies entirely with the author(s).*

*Founded in 1966, the American Society of Nephrology (ASN) is the world's largest professional society devoted to the study of kidney disease. Comprised of 11,000 physicians and scientists, ASN continues to promote expert patient care, to advance medical research, and to educate the renal community. ASN also informs policymakers about issues of importance to kidney doctors and their patients. ASN funds research, and through its world-renowned meetings and first-class publications, disseminates information and educational tools that empower physicians.*

###