

EMBARGOED FOR RELEASE until December 23, 2010 – 5:00 PM (EDT)

ASN Contact: Shari Leventhal • 202-416-0658 (p) • sleventhal@asn-online.org

KIDNEY DISEASE PATIENTS: EAT YOUR VEGGIES, REWARD YOUR KIDNEYS

Vegetarian Diet Lowers Blood and Urine Phosphorous Levels

Washington, DC (December 20, 2010) — Phosphorous levels plummet in kidney disease patients who stick to a vegetarian diet, according to a study appearing in an upcoming issue of the *Clinical Journal of the American Society Nephrology* (CJASN). The results suggest that eating vegetables rather than meat can help kidney disease patients avoid accumulating toxic levels of this mineral in their bodies.

Individuals with kidney disease cannot adequately rid the body of phosphorus, which is found in dietary proteins and is a common food additive. Kidney disease patients must limit their phosphorous intake, as high levels of the mineral can lead to heart disease and death. While medical guidelines recommend low phosphorus diets for patients with chronic kidney disease (CKD), phosphorus content is not listed on food labels.

Sharon Moe, MD (Indiana University School of Medicine and Roudebush Veterans' Affairs Medical Center) and her colleagues studied the effects of vegetarian and meat-based diets on phosphorous levels in nine patients with CKD. Patients followed a vegetarian or meat-based diet for one week, followed by the opposite diet two-to four-weeks later. Blood and urine tests were performed at the end of each week on both diets.

Despite equivalent protein and phosphorus concentrations in the two diets, patients had lower blood phosphorus levels and decreased phosphorus excretion in the urine when they were on the vegetarian diet compared with the meat-based diet. While the investigators did not determine the reason for this difference, a grain-based diet has a lower phosphate-to-protein ratio and much of the phosphate is in the form of phytate, which is not absorbed in humans.

The authors concluded that their study demonstrates that the source of protein in the diet has a significant effect on phosphorus levels in patients with CKD. Therefore, dietary counseling of patients with CKD must include information on not only the amount of phosphorous but also the source of protein from which it derives. "These results, if confirmed in longer studies, provide rationale for recommending a predominance of grain-based vegetarian sources of protein to patients with CKD. This diet would allow increased protein intake without adversely affecting phosphorus levels," the researchers

wrote.

Study co-authors include Miriam Zidehsarai, MD, Mary Chambers, RN, Laurie Trevino, MS (Indiana University School of Medicine); Lisa Jackman, MS, J. Scott Radcliffe, PhD (Purdue University); Susan Donahue, MBA, and John Asplin, MD (Litholink, Inc.)

Disclosures: Sharon Moe is a consultant and has received honoraria and/or grant support from Shire, Genzyme, and Ineos, makers of phosphate binders. She also serves as a Councilor for the American Society of Nephrology (ASN). Susan Donahue and John Asplin are employees of Litholink, which analyzed the urine studies.

The article, entitled “Vegetarian Compared with Meat Dietary Protein Source and Phosphorus Homeostasis in Chronic Kidney Disease,” will appear online at <http://cjasn.asnjournals.org/> on December 23, 2010, doi 10.2215/CJN.05040610.

The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.

Founded in 1966 and comprised of more than 12,000 members, the American Society of Nephrology (ASN) leads the fight against kidney disease by educating health professionals, sharing new knowledge, advancing research, and advocating the highest quality care for patients.

###