

EMBARGOED FOR RELEASE UNTIL OCTOBER 21 AT 9:00 AM ET

CONTACTS: Kurtis Pivert • kpivert@asn-online.org • 202-699-0238

Bob Henkel • bhenkel@asn-online.org • 202-557-8360

ASN KIDNEY WEEK 2014 • NOVEMBER 11-16 • PHILADELPHIA, PA

BINGE DRINKING IN YOUNG MEN LINKED WITH INCREASED RISK OF HYPERTENSION

Philadelphia, **PA** (October 21, 2014) — Binge drinking in early adulthood is associated with an increased likelihood of high blood pressure in males, while low to moderate alcohol use in early adulthood is associated with a decreased likelihood of hypertension in females. The findings come from a study that will be presented at ASN Kidney Week 2014 November 11–16 at the Pennsylvania Convention Center in Philadelphia, PA.

While studies have found that drinking alcohol can raise blood pressure in adults, little is known about the links between alcohol use during adolescence and hypertension. Researchers led by Sarah Twichell, MD (Boston Children's Hospital) analyzed data from the Growing Up Today Study (GUTS), a study of children who were 8 to 14 years old in 1996 and were followed with detailed surveys every 1 to 2 years. The team examined information on 8,605 participants who completed the 2010 survey.

Among the major findings:

- In young adult men, frequent binge drinking over the past year was associated with a 1.7-times increased likelihood of developing hypertension.
- In young adolescent males, there was no significant association between binge drinking or quantity of alcohol use and hypertension after they entered adulthood.
- In young women, binge drinking was not associated with hypertension.
- Light and moderate alcohol use in young adult women was associated with a significantly reduced likelihood of hypertension.

"Further study of alcohol use in young adulthood may provide insights into the early development of hypertension," said Dr. Twichell.

Study: "Adolescent Alcohol Use and the Development of Hypertension in Early Adulthood" (Abstract SA-PO156)

Disclosures: none.

ASN Kidney Week 2014, the largest nephrology meeting of its kind, will provide a forum for more than 13,000 professionals to discuss the latest findings in renal research and engage in

educational sessions related to advances in the care of patients with kidney and related disorders. Kidney Week 2014 will take place November 11–16, 2014, in Philadelphia, PA.

The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.

Founded in 1966, and with more than 15,000 members, the American Society of Nephrology (ASN) leads the fight against kidney disease by educating health professionals, sharing new knowledge, advancing research, and advocating the highest quality care for patients.

###