

EMBARGOED FOR RELEASE until May 7, 2015 – 5:00 PM (ET)

Contacts: Tracy Hampton • (312) 339-9067 • thampton@nasw.org
Bob Henkel • (202) 557-8360 • bhenkel@asn-online.org

DEXAMETHASONE MAY HELP PREVENT SEVERE KIDNEY INJURY FOLLOWING HEART SURGERY

The inexpensive anti-inflammatory drug could save thousands of lives each year

Highlights

- Patients who received dexamethasone during heart surgery had about a 2.5-times lower risk of developing kidney failure requiring dialysis compared with those receiving a placebo.
- The greatest benefits of dexamethasone were seen in patients with pre-existing advanced chronic kidney disease.

Acute kidney injury can be a serious complication following heart surgery.

Washington, DC (May 7, 2015) — The anti-inflammatory drug dexamethasone helps prevent serious kidney complications that can arise following heart surgery, according to the results of a randomized clinical trial. The findings, which appear in an upcoming issue of the *Journal of the American Society of Nephrology* (JASN), could lead to a change in care for patients during cardiac operations.

Acute kidney injury (AKI) is one of the most devastating complications following cardiac surgery. Approximately 1% of patients undergoing cardiac surgery require dialysis to treat severe AKI that arises after surgery, and the incidence is higher among patients with pre-operative chronic kidney disease. These patients experience strikingly high death rates while in the hospital that exceed 40%. “One percent sounds like a small percentage, however given the fact that each year, over half a million people undergo heart surgery in the United States alone, this means that an estimated 5000 patients develop renal failure and of those about 2500 die as a result of this complication,” said Kirolos Jacob, MD (University Medical Center, Utrecht, The Netherlands). He noted that these figures are rising due to the aging population.

Because heart surgery initiates an inflammatory reaction in the body that can have negative effects on the kidneys, Dr. Jacob and his colleagues wondered whether giving patients dexamethasone, an anti-inflammatory drug, could decrease the risk of severe AKI following cardiac surgery. The team analyzed the results of a large randomized controlled trial called the Dutch Dexamethasone for Cardiac Surgery (DECS) trial, which included 4465 patients undergoing cardiac surgery who were randomized to receive

placebo or dexamethasone during surgery. The original trial tested whether dexamethasone could reduce the risk of a variety of major postoperative complications. In this analysis, the investigators specifically examined kidney failure and focused on the most severe form: AKI requiring dialysis.

Dexamethasone appeared to protect against the development of severe AKI. Patients who received the drug had about a 2.5-times lower risk of developing AKI requiring dialysis compared with those receiving a placebo.

“The beneficial effects of dexamethasone were particularly present in those who already had pre-existing kidney disease before heart surgery,” said Dr. Jacob. “This reinforces the fact that this drug could be of major importance for the increasing elderly population with pre-existing kidney disease undergoing a heart operation.”

The study is the largest randomized, placebo-controlled trial showing a potential benefit of any therapeutic drug for the prevention of severe kidney injury following heart surgery. A single dose of dexamethasone during a heart operation is inexpensive, straightforward, painless, and safe for patients. “These advantages make the intervention very accessible and cost-effective, especially since the costs for dialysis are very high,” said Dr. Jacob.

Study co-authors include David Leaf, MD, MMSc, Jan Dieleman, MD, Diederik van Dijk, MD, PhD, Arno Nierich, MD, PhD, Peter Rosseel, MD, PhD, Joost van der Maaten, MD, PhD, Jan Hofland, MD, PhD, Jan Diephuis, MD, Fellery de Lange, MD, PhD, Christine Boer, PhD, Jolanda Kluin, MD, PhD, Sushrut Waikar, MD, MPH, for the Dexamethasone for Cardiac Surgery (DECS) Study Group.

Disclosures: The DECS study was sponsored by grants from the Netherlands Organisation for Health Research and Development (ZonMw) and the Dutch Heart Foundation.

The article, entitled “Intraoperative High-Dose Dexamethasone and Severe Acute Kidney Injury after Cardiac Surgery,” will appear online at <http://jasn.asnjournals.org/> on May 7, 2015.

The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.

The American Society of Nephrology®, ASN®, Kidney Week®, CJASN®, JASN®, NephSAP®, and ASN Kidney News® are registered trademarks of ASN

Founded in 1966, and with more than 15,000 members, the American Society of Nephrology (ASN) leads the fight against kidney disease by educating health professionals, sharing new knowledge, advancing research, and advocating the highest quality care for patients.

#

Tweet: **Dexamethasone helps prevent severe kidney injury following heart surgery.**
<http://www.bit.ly/ASN-XXXX>

Facebook: The anti-inflammatory drug dexamethasone helps prevent serious kidney complications that can arise following heart surgery, according to the results of a randomized clinical trial. The findings, which appear in the *Journal of the American Society of Nephrology*, could lead to a change in care for patients during cardiac operations..

&&

Head Coordinator of the Communications and Media Division coordinator of the University Medical Center in Utrecht, the Netherlands:

Mr. Paul Geurts
University Medical Center Utrecht
Room number: C01.212
Internal post mail: D01.343
Post box 85500
3508 GA UTRECHT
Tel: +31 88 75 551 90
E-mail: T.B.P.Geurts@umcutrecht.nl

Brigham and Women's Health Media Relations Division:
bwhmediarelations@partners.org