

**EMBARGOED FOR RELEASE until September 17, 2015 – 5:00 PM (ET)**

**Contacts:** Tracy Hampton • (312) 339-9067 • [thampton@nasw.org](mailto:thampton@nasw.org)  
Bob Henkel • (202) 557-8360 • [bhenkel@asn-online.org](mailto:bhenkel@asn-online.org)

## **HIGH DIETARY SODIUM AND POTASSIUM MAY WORSEN CHRONIC KIDNEY DISEASE**

*Reducing intake may safeguard patients' health*

### **Highlights**

- High urinary excretion levels of both sodium and potassium were linked with faster progression of chronic kidney disease.
- Patients with chronic kidney disease tend to consume sodium above the recommended daily limit.

*An estimated 26 million people in the United States have chronic kidney disease.*

**Washington, DC (September 17, 2015)** — High dietary intake of sodium and potassium may speed the progression of kidney disease, according to a study appearing in an upcoming issue of the *Journal of the American Society of Nephrology (JASN)*. The findings could impact dietary recommendations to help safeguard patients' health.

Chronic kidney disease (CKD) is a major public health challenge because it is common, frequently progresses to kidney failure, and increases risk of heart disease and premature death. Diet may play an important role in CKD progression, but little is known about the role of certain dietary components such as sodium and potassium.

To investigate, Jiang He, MD, PhD (Tulane University) and his colleagues collected yearly urine samples from 3939 patient with CKD to estimate the patients' dietary sodium and potassium intake. The researchers found that high urinary excretion levels of both sodium and potassium were linked with faster disease progression. In addition, the study's participants consumed an average of 3700 mg of sodium per day, which is much higher than the recommended 2400 mg per day limit.

"These data warrant future clinical trials to test the effect of a moderate reduction in dietary sodium and potassium intake on CKD progression in patients with high dietary sodium or potassium intake," said Dr. He. "The findings could ultimately impact dietary recommendations for patients with CKD to slow disease progression."

Study co-authors include Katherine Mills, PhD, Lawrence Appel, MD, MPH, Wei Yang, PhD, Jing Chen, MD, MS, Belinda Lee, MD, Sylvia Rosas, MD, Anna Porter, MD, Gail Makos, RN, MSN, Matthew Weir, MD, L. Lee Hamm, MD, and John Kusek, PhD.

Disclosures: The authors reported no financial disclosures. The research was funded by the National Institute of Diabetes and Digestive and Kidney Diseases, grant number R01-DK074615.

The article, entitled “Urinary Sodium and Potassium Excretion and Chronic Kidney Disease Progression,” will appear online at <http://jasn.asnjournals.org/> on September 17, 2015.

*The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.*

*Founded in 1966, and with more than 15,000 members, the American Society of Nephrology (ASN) leads the fight against kidney disease by educating health professionals, sharing new knowledge, advancing research, and advocating the highest quality care for patients.*

# # #

Tweet: High dietary sodium and potassium may worsen chronic kidney disease.  
<http://www.bit.ly/ASN-XXXX>. Author’s Twitter handle:

Facebook: High dietary intake of sodium and potassium may speed the progression of kidney disease, according to a study in the *Journal of the American Society of Nephrology*. The findings could impact dietary recommendations to help safeguard patients’ health.

\*&\*&

Keith Brannon, Associate Director of Public Relations  
[kbrannon@tulane.edu](mailto:kbrannon@tulane.edu)  
504-862-8789

The American Society of Nephrology®, ASN®, Kidney Week®, CJASN®, JASN®, NephSAP®, and ASN Kidney News® are registered trademarks of ASN